

Quantum Horizons and Space-Time Non-Commutativity

M. Martinis and V. Mikuta-Martinis

Rudjer Bošković Institute, Zagreb, Croatia

Abstract

We study dynamics of a scalar field in the near-horizon region described by a static Klein-Gordon operator which is the Hamiltonian of the system. The explicit construction of a time operator near-horizon is given and its self-adjointness discussed.

* Expanded version of a poster presented at the 5 European Advanced Study Conference in Ancient Olympia, Greece, May 2004.

1 Introduction

Einstein's general theory of relativity [Nicolson,1981] describes gravity as a curvature of spacetime caused by the presence of matter. A black hole is formed if a massive object (e.g. a star) collapses into an infinitely dense state known as a singularity. In this case, the curvature of spacetime becomes extreme and prevents any light or other electromagnetic radiation from escaping to infinity. The event horizon defines the boundary of a black hole behind which nothing, not even light, can escape.

Applying the Einstein Field Equations to collapsing stars, German astrophysicist Kurt Schwarzschild introduced the critical radius $R_S = 2GMc^{-2}$ for a given mass at which matter would collapse into a singularity. The simplest three-dimensional geometry for a black hole is a sphere (known as a Schwarzschild black hole), its surface defines the event horizon.

Therefore, in the standard General Relativity without Quantum Mechanics (QM) a fundamental instability against collapse implies the existence of black holes as stable solutions of Einstein's equations. The QM and the Quantum Field Theory (QFT) in the curved spacetime with classical event horizon are, however, troubled by the singularity at the horizon [t Hooft, 2004]. This problem may be solved by treating the black hole as a quantum state which implies that the energy of the black hole and its corresponding time do not commute at the horizon [Bai and Yan, 2004]. In this picture we study the complex dynamics of a scalar field in the near-horizon region described by a static Klein-Gordon operator which in this case becomes the Hamiltonian of the system. We present the explicit construction of the time operator near-horizon and discuss its self-adjointness [Martinis and Mikuta,2003].

2 Scalar Field in the Near-Horizon Region

The Schwarzschild geometry of a static spherical black hole is described by the metric

$$ds^2 = -f(r)dt^2 + [f(r)]^{-1}dr^2 + r^2d\Omega^2, \quad (1)$$

where $rf(r) = r - 2M$ and $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$. This metric is minimally coupled to a scalar field $\Phi(r, t, \Omega)$ through the action ($c = 1, \hbar = 1$ and $G = 1$):

$$S = -\frac{1}{2} \int d^4x \sqrt{-g} [g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi + m^2 \Phi^2], \quad (2)$$

where $g = \det[g^{\mu\nu}]$. The equation of motion in the black-hole gravitational background is

$$\begin{aligned} (\square_g - m^2)\Phi &= \frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\Phi) - m^2\Phi \\ &= -\frac{1}{f}\Phi_{tt} + f\Phi_{rr} + (f' + 2f/r)\Phi_r + \triangle_\Omega\Phi/r^2 - m^2\Phi \\ &= 0. \end{aligned} \quad (3)$$

By separating the time and angular variables

$$\Phi(t, r, \Omega) = e^{-it\omega} f^{-1/2} r^{-1} \psi(r) Y_{lm}(\Omega), \quad (4)$$

the behavior of the Schroedinger-like equation near the horizon can be studied by means of an expansion in the near-horizon variable $x = r - 2M$. In this variable, for small x , we obtain a conformally invariant Schroedinger equation

$$\left[\frac{d^2}{dx^2} + \left(\frac{1}{4} + \Theta^2\right)x^{-2}\right]\psi(x) = 0 \quad (5)$$

where $\Theta = 2M\omega$. This reduction to an effective Schroedinger-like equation can be described by an effective attractive potential $V(x) = -(1/4 + \Theta^2)x^{-2}$ which is conformally invariant with respect to the near-horizon variable x . The corresponding quantum Hamiltonian is

$$H = (p^2 + \frac{g}{x^2}), \quad (6)$$

where $g = -2(1/4 + \Theta^2)$ is supercritical for all nonzero frequencies ω .

3 Time Operator and Spacetime Noncommutativity

The singularity associated with horizon may be overcome by assuming space-time non-commutativity on the horizon, the so called Quantum Horizon [t Hooft, 2004], [Bai and Yan, 2004]. If a black hole is considered as a quantum state, its energy $\hat{H}$ and its conjugate time $\hat{t}$ are expected to become conjugate operators obeying

$$[\hat{H}, \hat{t}] = i. \quad (7)$$

Since $\hat{H}$ is x-coordinate dependent, we expect spacetime noncommutativity, $[\hat{t}, x] \neq 0$. What is $\hat{t}$ as an operator? Due to Pauli theorem [Pauli,1958] no such self-adjoint operator should exist if the spectrum of the Hamiltonian is semibounded or discrete. In quantum theory $\hat{H}$ has a continuous spectrum for $g \geq 3/4$ with $E > 0$ but no ground state at $E = 0$. It is essentially self-adjoint only for $g > 3/4$ in the domain

$$\mathcal{D}_0 = \{\psi \in \mathcal{L}^2(dx), \psi(0) = \psi'(0) = 0\} \quad (8)$$

For $g \leq 3/4$, [Birmingham *et al.*, 2001] the Hamiltonian is not essentially self-adjoint, but it admits a one-parameter family of self-adjoint extension labeled by a $U(1)$ parameter e^{iz} , where z is real, which labels the domains $\mathcal{D}_z$ of the extended Hamiltonian. The set $\mathcal{D}_z$ contains all the vectors in $\mathcal{D}_0$, and vectors of the form $\psi_+ + e^{iz}\psi_-$.

For $g = -\frac{1}{4}$, and a given self-adjoint extension z , there are infinite number of bound states:

$$\psi_n(x) = N_n \sqrt{x} K_0 \left(\sqrt{E_n} x \right), \quad (9)$$

$$E_n = \exp \left[\frac{\pi}{2} (1 - 8n) \cot \frac{z}{2} \right], \quad (10)$$

where n is an integer, N_n is a normalization factor, and K_0 is the modified Bessel function. The horizon, in this picture, is located at $x = 0$ where the wave function ψ_n vanishes. However, in a band-like region $\Delta = [x_0(1 - \delta), x_0(1 + \delta)]$, where $x_0 \sim E_0^{-1/2}$ with $z > 0$, all the eigenfunctions of $\hat{H}$ exhibit a scaling behaviour of the type, $\psi_n \sim \sqrt{x}$.

Formally, the three generators: $\hat{H}$, the scaling operator $\hat{D} = -(xp + px)/4$ and $\hat{K} = x^2/2$ obey the conformal algebra commutation relations

$$[\hat{H}, \hat{D}] = i\hat{H}, \quad [\hat{K}, \hat{D}] = -i\hat{K}, \quad [\hat{H}, \hat{K}] = 2i\hat{D}, \quad (11)$$

with a constant quadratic Casimir operator

$$\begin{aligned} C_2 &= \frac{1}{2}(\hat{K}\hat{H} + \hat{H}\hat{K}) - \hat{D}^2 \\ &= \frac{g}{4} - \frac{3}{16}. \end{aligned} \quad (12)$$

If $\hat{H}^{-1}$ exists [Klauder,1999], we can formally construct a time operator $\hat{t}$ [Martinis and Mikuta,2003]

$$\hat{t} = \frac{1}{2}(\hat{D}\hat{H}^{-1} + \hat{H}^{-1}\hat{D}), \quad (13)$$

which obeys the required commutation relation, $[\hat{H}, \hat{t}] = i$. Although, both $\hat{H}$ and $\hat{D}$ separately can be made self-adjoint operators it is not true for a $\hat{t}$ -operator which contains $\hat{H}^{-1}$ [Klauder,1999]. In the limit $g \rightarrow 0$, we have $\hat{H} \rightarrow H_0 = p^2/2$ and $\hat{t} \rightarrow t_0$ where

$$\hat{t}_0 = -\frac{1}{2}(xp^{-1} + p^{-1}x) \quad (14)$$

is exactly the time-of-arrival operator of Aharonov and Bohm [1961] The Aharonov-Bohm time operator $\hat{t}_0$ is not self-adjoint and its eigenfunctions are not orthonormal.

4 Conclusion

In this paper, we have studied the properties of a scalar field in the near-horizon region of a massive Schwarzschild black hole. The quantum Hamiltonian governing the near-horizon dynamics is found to be scale invariant and has the full conformal group as a dynamical symmetry group. Using only the generators of the conformal group, we constructed the time operator near-horizon. The self-adjointness of $\hat{H}$ and $\hat{t}$ is also discussed.

References

- Aharonov, Y. & Bohm, D. [1961] "Time in the Quantum Theory and the Uncertainty Relation for Time and Energy," *Phys. Rev.* **122**, 1649- .
- Bai, H. & Yan, Mu-Lin [2004] *Quantum Horizon*, (gr-qc/0403064).
- Birmingham, D. & Gupta, K. S. & Sen, S. [2001] "Near-Horizon Conformal Structure of Black Holes," *Phys. Lett.* **B505**, 191-196. 't Hooft, G. [2004] *Horizons* (gr-qc/0401027).
- Klauder, J.R. [1999] *Noncanonical Quantization of Gravity I. Foundation of Affine Quantum Gravity*, (gr-qc/9906013).
- Martinis, M. & Mikuta, V. [2003] "Time Operator for a Singular Quantum Oscillator" *Fizika* **B12**, 285-290.
- Nicolson, I. [1981] *Gravity, Black Holes, and the Universe* (David and Charles, London).
- Pauli, W. [1958] "Die allgemeinen Prinzipien der Wellenmechanik" in *Encyclopedia of Physics*, ed. Fluegge, S. (Springer-Verlag, Berlin) pp. 1-168.